

*******PLEASE REVIEW ENTIRE PAGE TO AVOID POSSIBLE CANCELLATION/RESCHEDULE OF PROCEDURE*******

Please notify us if you take any of the following medications:

Coumadin (Warfarin), Plavix (clopidogrel), Aggrenox (dipyridamole ER), Persantine (dipyridamole), Ticlid (ticlopidine), Pletal (cilostazol), Effient (prasugrel), Xarelto (rivaroxaban), Brilinta (ticagrelor), Pradaxa (dabigatran), or Eliquis (apixaban).

HOLD THE FOLLOWING MEDS 7 DAYS PRIOR TO THE PROCEDURE:

PLEASE NOTE There are often times a medication will have a different name. If you are concerned about needing to hold a medication you are taking please notify our office at 918-207-0025

Advil	Congesprin	Gelprin	Pinstel
Aleve	Cope	Ibuprofen	Pradaxa
Alka Seltzer	Coricidin	Indocin	Relafen
Anacin	Coumadin	Indomethacin	Robaxin
Anaprox	Damason	Ketoprofen	Salfelx
Ansaid	Darvon with ASA	Lortab ASA	Sine-Aid
APC	Daypro	Meclomen	Sine-Off
Arthrotec	Disalcid	Medipren	Soma CMD
Ascodeen	Dolobid	Meloxicam	Sulindac
Ascriptin	Dristan	Midol	Tolectin
Axdone	Easprin	Motrin	Toradol
BC Powder	Exotrin	Nalfon	Trigesic
Bufferin	Empirin	Naprosyn	Trilisate
Bruden	Emprazil	Norgesic	Tylenol Arthritis
Cataflam	Equagesic	Nuprin	Vanquish
Celebrex	Excederin Caps	Orphengesic	Vitamin E
Cephalgesic	Feldene	Orudis	Voltaren
Cheracol	Florinal	Percodan	Zactin
Clinoril	Four way cold tabs	Piroxican	Phenaphen

******* IF YOUR PHYSICIAN HAS PLACED YOU ON LOW DOSE OR REGULAR STRENGTH ASPIRIN FOR A SPECIFIC PROBLEM/ILLNESS PLEASE CONTINUE TO TAKE IT, FOR PREVENTATIVE PURPOSES PLEASE STOP 7 DAYS PRIOR TO THE PROCEDURE. *******

If you need minor pain medication you may take Acetaminophen (Tylenol) or another non-aspirin medication. Tylenol or Anacin-3 are available at your local pharmacy without a prescription and have comparable pain relief potential to that of aspirin.